



ITALPOLLINA
HELLO NATURE!

MYR

CALCIUM - BORON




LEAF LIQUID INTEGRATORS

MYR RANGE is based on micronutrients and meso elements complexed with vegetal peptides with the aim of prevent and/or cure nutritional deficiencies and improve yield and fruit quality.

MYR RANGE guarantees the highest efficiency and a quick absorption, without losses due to insolubilization. It is the result of a careful research on the mechanisms of vegetal absorption in order to prevent phenomena of phytotoxicity caused by micronutrients.

The peptides of **MYR** are quickly absorbed and then incorporated in the lymph : they can carry out their functions in stressed plants in a few hours, allowing a remarkable saving of energy and permitting the recovery of the optimal conditions for development.

Oligonutritional complexes **MYR** have a stable structure also in alkaline environment, which prevents the formation of salts or insoluble hydroxides.



1) Nutrients arrive on leaves

2) Foliar absorption

3) Translocation of the nutrients inside the plants.

MYR RANGE is powered by **LISIVEG**, an exclusive production system designed by the R&D of

LISIVEG® *Italpollina*, based on the **enzymatic hydrolysis of vegetal proteins**. It allows to obtain **PEPTIDES** that, unlike individual free amino acids, have a more prominent, assured and **natural biostimulant effect**. **LISIVEG**® lies its strength in the innovative treatment technology that **preserves even the weakest components** (as Tryptophan) and maintains its activity unaltered.



MYR CALCIUM - BORON

MYR CALCIUM - BORON improves plant growth and development and has an anti-stress and nutritional effect. Its action is fast and effective.

Organic Nitrogen (N)	3%
Calcium oxide (CaO) water soluble	5%
Boron (B) water soluble	1%
Organic carbon (C)	19%
pH (1:5)	6
Specific weight	1,23 Kg/L

THE ADVANTAGES

- Prevention of sensitive calcium physiological disorders
- Improve of formation, quality and shelf life of fruits
- Increase the tolerance of plants to environmental stresses
- Increase the pollen fertility
- Improvement of photosynthesis and fruit setting, reducing fruit drop
- Quick nutritional action



USE & DOSE

All dosages provided are in standard conditions, dosages and modes of application are to be determined on the basis of the requirements and the nutritional status of the crop.

We recommend the application in the coolest hours of the day (morning or evening) to obtain maximum absorption by the leaves and the best effectiveness.

MYR range can be mixed with the most common phytosanitary products and fertilizers on the market. Before using we recommend testing on a small surface for compatibility with other products.

Liquid fertilizer for foliar application (on leaves).

Crops	Dose recommended		Stage
	Dosage per ha	Dosage per hl	
Vegetables	3L	300mL	Every 15 days
Orchards	3L	300mL	Before flowering period - 2-3 applications
Soft fruits	2L	300mL	Every 15 days starting in preflowering period
Vineyards, ananas, bananas, kiwi	3L	300mL	Before flowering period - 2-3 applications
Sunflower	3L	1,5L*	Limit tractor crossing 7-8 leaves
Sugar beet	3L	1,5L*	2 applications to cover crop needs

*based on 200 L/ha of spray solution

MYR

CALCIUM - BORON



PACKAGING
1 - 5 - 20 L



*Suitable for organic farming
according to EC REG. 834/2007*

ITALPOLLINA S.p.A.
Località Casalmenini, 10
37010 Rivoli Veronese (VR) Italia
Tel +39.045.6205511
info@italpollina.com
www.italpollina.com

Follow us   